

# **FORT BERTHOLD RESERVATION**

## **List of Topics**



### **BACKGROUND**

Reservation Overview

Production Overview

### **GEOLOGIC OVERVIEW**

Geologic History

Petroleum Systems

Summary of Play Types

### **CONVENTIONAL PLAY TYPES**

Play 1 - Folded Structure-Mississippian Carbonate Play

Play 2 - Mississippian Shoreline Play

Play 3 - Mississippian Lodgepole Waulsortian Mounds

Play 4 - Ordovician Red River Play

Play 5 - Devonian Nisku-Duperow Play

Play 6 - Pre-Prairie (Winnipegosis/Interlake Play)

Play 7 - Post Madison Clastics (Tyler-Heath)

Play 8 - Pre-Red River Gas Play

Play 9 - Bakken Fairway/Sanish Sand Play

### **UNCONVENTIONAL / HYPOTHETICAL PLAY TYPES**

Play 10 - Niobrara Microbial Gas Play

### **REFERENCES**

# OVERVIEW

## FORT BERTHOLD RESERVATION

### The Three Affiliated Tribes

Tribal Headquarter: New Town, North Dakota  
 Geologic Setting: Williston Basin

### Introduction

The Fort Berthold Indian Reservation is located in west-central North Dakota approximately thirty miles southwest of the city of Minot. The Reservation contains portions of Dunn, McKenzie, McLean, Mercer, Mountrail and Ward Counties and includes an area of about 1,530 square miles or 980,000 acres. These lands are located 15 miles east of the center of the Williston Basin, a geologic area where undiscovered accumulations of oil and gas may be located.

Several studies have been published over the years which indicate high potential for undiscovered oil and gas reserves on the Fort Berthold Reservation. There has been past interest exhibited by oil companies, however, high royalties, high lease acquisition costs, inability to assemble large blocks of acreage, rights to seismic data, Tribal Employment Rights Office (TERO) regulations, taxes, and a 100 percent signature requirement imposed by the federal statute on Trust lands have served as deterrents to oil and gas exploration on the Reservation. The 100 percent signature requirement regulation has made exploration on Tribal Allotted Lands nearly impossible to carry out due to the high fragmentation caused by heirship. The Tribes are currently working to correct these problems to open the door for future gas and oil exploration and development. The Three Affiliated Tribes are striving to work closer with oil companies to make oil and gas exploration on Fort Berthold competitive with lands outside of the Reservation.

The Fort Berthold Reservation possesses all the requisites for commercial petroleum development. According to an oil and gas study authorized by Joe H. Rawlings, source rocks and reservoir caprock combinations are in evidence from the Antelope field located near the northwest corner of the Reservation. This field produces both oil and gas from four different zones. The relatively new Plaza field, located near the east exterior boundary, is also a major producing oil field. Other fields were recently discovered on the Reservation while drilling the Bakken and Mission Canyon formations. This multiplicity of geologic structures argues the presence of the many deep traps. Regardless of the development of these fields, much of the Reservation has not been explored for accumulations of oil and gas.

The Williston Basin, which encompasses the Reservation, has a long history of production. Much of the oil in this area was sourced by the organically rich Bakken Formation. New horizontal drilling technology has made production from Bakken source rocks possible. A report written for the Bureau of Indian Affairs by Susan Race Wager, states that the Fort Berthold Reservation is favorably located for exploration in the Bakken Formation.

According to George Long of the Bureau of Land Management (BLM), there have been 571 tests for oil and gas on or immediately adjacent to the Fort Berthold Reservation resulting in a total of 392 producing wells and 179 plugged and abandoned wells; a 69% success ratio. The majority of these 179 plugged and abandoned wells did report oil and gas shows. Of special interest, there appears to be production potential in the Mississippian Charles Formation which has been bypassed in all of the wells drilled except those few that are too shallow to reach the Charles Formation located in the Mission Canyon.

Water saturation calculations were made on 60 wells to evaluate possible bypassed production in the Mississippian Charles Formation. Possible oil and gas production was indicated in 52 of these 60 wells. According to the BLM,

there are approximately 10 formations proved to be productive in the Fort Berthold area. Of further note, the facies distribution during lower Mississippian time strongly suggests that Lodgepole trends are present on the Fort Berthold Indian Reservation (USGS).

The Three Affiliated Tribes have purchased seismic data from lines located in the western portion of the Reservation, which may be examined by parties interested in oil and gas exploration. Some of the seismic data will be reprocessed and may be correlated with borehole logs. Sections and data tapes reside with the Division of Energy and Minerals Resources of the Bureau of Indian Affairs located in Denver, CO.

### Area Location and Access

Fort Berthold Indian Reservation comprises parts of Dunn, McKenzie, McLean, Mercer, Mountrail, and Ward Counties in west-central North Dakota (Figure 1), near the confluence of the Missouri and Little Missouri River valleys. Total area is about 1,530 square miles, approximately 11 percent of which is covered by waters impounded by Garrison Dam (Lake Sakakawea). The lake divides the reservation into four distinct areas, here referred to as the western, southern, eastern, and north-central segments.

Although reservoir waters somewhat impede travel between the four land segments, most of the reservation is accessible over a system of State highways and local roads. Rail service is provided to the northern part of the reservation by the Soo Line Railroad. A main east-west line of the Burlington Northern passes within 7 miles of the reservation, roughly paralleling the southern boundary.

### Physiography

The Fort Berthold Indian Reservation includes land that ranges from rugged badlands to rolling plains. Altitudes range from about 1,850 feet at Lake Sakakawea to over 2,600 feet on Phaelen's Butte near Mandaree. The reservation is within the Northern Great Plains Physiographic Province and may be divided into four physiographic units: (1) the Coteau Slope; (2) the Missouri River trench (now flooded); (3) the Missouri Plateau; and (4) the Little Missouri Badlands. South of Lake Sakakawea the reservation has a bedrock surface with scattered areas of glacial drift. North of the lake, glacial deposits predominate and only patches of bedrock crop out. The landscape reflects this distribution of sediments: south of the lake, hills and badlands are common; north of the lake the glaciated topography is mainly undulating to rolling.

The reservation area north of Lake Sakakawea is part of the Coteau Slope, which has both erosional and glacial landforms with glacial predominating. Gentle slopes characterize 50 to 80 percent of the area and local relief ranges from 50 to 200 feet. The Little Missouri Badlands lie adjacent to the Little Missouri River south and west of Lake Sakakawea as well as in a few restricted areas along the Missouri River. They consist of rugged, deeply-eroded, hilly land in which gentle slopes characterize only 20 to 50 percent of the area and local relief is commonly over 500 feet. Areas other than badlands south and west of the lake are part of the Missouri Plateau. In these areas, gentle slopes characterize about 50 to 70 percent of the area and local relief ranges from 300 to 500 feet.

The Missouri and Little Missouri Rivers and their larger tributaries have cut deeply into the bedrock and glacial deposits of various compositions. The Missouri River is 300 to 500 feet below the upland plain. Near the western boundary of the reservation, the Little Missouri River has eroded a channel more than 600 feet deep. Occasional ridges and bare buttes extend as much as 400 feet above the plain.

### Land Status

The Fort Berthold Indian Reservation was established by the Fort Laramie Treaty of September 17, 1851, for the Arikara, Mandan, and Hidatsa Tribes of Indians who later united to form the Three Affiliated Tribes. Executive Orders and Congressional Acts have limited the reservation to its present boundaries. The act of June 1, 1910, 36 Stat. 455, opened unallotted and unsold reservation lands to non Indians, thus creating the "ceded and diminished lands" boundary. It was assumed by many that only the remaining lands comprised the Fort Berthold Indian Reservation. A Federal appeals court (8th Cir. 1972), however, ruled that the 1910 Act did not change reservation boundaries and that the "homestead" (ceded) area remained a part of the reservation (City of New Town vs. United States, 454 F 2d 121) Public Law 437 and the Act of July 31, 1947 (amended October 29, 1947) made provision for lands inundated by the Garrison Dam reservoir. Table 1 summarizes the present extent of land holdings on the Fort Berthold Indian Reservation. Most of the north and northeast part of the reservation (the homestead area) is in private ownership. Land status data are from Bureau of Indian Affairs records.


Nearly 54 percent of the reservation's subsurface mineral rights are owned by the Three Affiliated Tribes. Mineral rights in the diminished reservation area are all tribally owned with the exception of 164.09 acres owned by the Federal government. The Tribes also retain mineral ownership for 110,623.13 acres of the homestead area. Lands in the Garrison reservoir area were severed.

**TABLE 1** - Summary of land ownership, Fort Berthold Indian Reservation, N. Dakota

Classification	Acreage	Percentage of total
<b>Diminished Reservation Area</b>		
Tribally-owned lands.	57,954.20	5.91
Allotted lands	360,438.57	36.76
Government-owned land	164.09	0.01
Privately owned (alienated) land	55,865.14	5.70
Subtotal	474,422.00	48.38
Reservoir Taking Area	152,359.95	15.54
Homestead (ceded) Area.	353,792.59	36.08
Total area of reservation.	980,574.54	100.00

### Contacts

Inquiries concerning oil and gas leases on the Fort Berthold Reservation may be directed to the Three Affiliated Tribes Natural Resources Department - telephone (701) 627-3627 or the Bureau of Indian Affairs, located in New Town, North Dakota - (701) 627-3741.



# INTRODUCTION

## Fort Berthold Reservation

### Williston Basin

Over 700 MMBO have been produced from the Williston Basin, one of the largest cratonic basins in North America. The reservation is ideally situated for numerous exploration targets within this basin. Several source rock horizons, including the world renowned Bakken Formation, contribute to the prolific nature of the basin.

The Williston Basin contains an estimated mean value of 650 MMBO and 1.69 TCFG from undiscovered resources in conventional plays. Multiple episodes of maturation and migration occurred during Permian-Cretaceous time from these source intervals. Understanding the trapping mechanisms and migration pathways are critical to successful future exploration within the reservation area. Carbonate reservoirs in Paleozoic formations have been the primary focus of hydrocarbon exploration. Recent exploration targets include microbial gas in Cretaceous sediments and deep Paleozoic sandstone intervals.

### Early Exploration in the Williston Basin and Fort Berthold Reservation

Early discoveries were made on large surface structures such as Nesson and Cedar Creek Anticlines, and Poplar Dome. The Williston Basin is distinctive among other Rocky Mountain basins because of its continuous basin subsidence and burial history throughout Paleozoic and Mesozoic time. Large volumes of clastic and carbonate sediments have been preserved.

Since the late 1940's, industry has found more than 960 fields and the basin has undergone multiple exploration cycles. The Williston Basin covers more than 143,000 sq. miles and Fort Berthold reservation covers about one percent of that total (1530 sq. miles). Most of the reservation is unexplored.

Gas was discovered at Cedar Creek Anticline in 1916; oil was discovered on Nesson Anticline in 1951. Nesson is located about 50 miles northwest of the reservation boundary. Antelope Field, a southeast plunging anticline, was discovered in 1953 and extends onto the reservation. Plaza and Wabek Fields, part of the Mississippian shoreline trend, were discovered in the 1980's. Condensate and gas were discovered in the Winnipeg and Deadwood Formations at Antelope Field in 1992.

### Fort Berthold Reservation GENERAL PRODUCTION INFORMATION

#### U.S.G.S. Geologic Province - Williston Basin (031) Tectonic Province - Williston Basin

#### Fields within reservation boundaries

1996 cumulative production. Parentheses indicates discovery year

- (1953) **Antelope** - 41 MMBO, 19.2 Mmcf, 30 oil wells, 2 gas wells
- (1989) **Plaza** - 2.9 MMBO, 3.9 Mmcf, 20 wells total
- (1982) **Wabek** - 5.4 MMBO, 3.9 Mmcf, 18 wells total

#### Nearby fields

- (1955) **Blue Buttes** - 46 MMBO, 29.2 Mmcf, 44 wells total
- (1957) **Bear Den** - 1.5 MMBO, 1.7 Mmcf, 2 oil wells, 1 gas well
- (1952) **Croff** - 1.8 MMBO, 4.1 Mmcf, 3 wells total
- (1981) **Spotted Horn** - 108 MBO, 36,234 Mcf, (Abn'd)
- (1982) **Squaw Creek** - 195 MBO, 328,546 Mcf, 1 well total
- (1982) **Mandaree** - 160 MBO, 147,325 Mcf, 2 wells total
- (1990) **Lucky Mound** - 1.4 MMBO, 890, 670 MCF, 18 wells total

**Figure FB-1.1.** Producing horizon legend. Many of the potential reservoir intervals can be correlated into Wyoming and Montana. However, the Williston Basin is unique among other Rocky Mountain basins for its thick package of Paleozoic age carbonate sediments. While the other basins are known for their numerous clastic potential reservoir intervals, the Williston Basin is known as a carbonate province (modified after Seventh International Williston Basin Symposium Guidebook, 1995).

PRODUCING HORIZON LEGEND							
ERA	SYSTEM	SERIES		WILLISTON BASIN	POWDER RIVER BASIN	WESTERN WYOMING SOUTHERN MONTANA	WESTERN NORTHERN MONTANA
CENOZOIC	TERTIARY			Fort Union	White River Wasatch Fort Union	Green River Wind River Wasatch Fort Union	Fort Union
				Fox Hills Judith River Eagle Niobrara Greenhorn	Lance Teckla Mesaverde Teapot Parkman Sussex Shannon Niobrara Frontier	Lance Fox Hills Mesaverde Cody Shannon Niobrara Frontier	Hell Creek Judith River Clagget Eagle Telegraph Creek Niobrara Greenhorn Frontier
MESOZOIC	CRETACEOUS	UPPER		Dakota Group	Mowry Muddy Dakota Fall River Lakota	Mowry Muddy Bear River Dakota Cloverly	Blackleaf Bow Island Kootenai Cat Creek Moulton Sunburst Cut Bank
		LOWER					
MESOZOIC	JURASSIC			Morrison Ellis Group Swift Reirdon Piper Nesson	Morrison Sundance Canyon Springs Gypsum	Gannet Morrison Sundance Stump-Preuss Twin Creek	Morrison Ellis Group Swift Reirdon Sawtooth
	TRIASSIC			Spearfish	Minnekahta Chugwater Spearfish	Nugget Chugwater Ankareh Thaynes Woodside	
	PERMIAN			Opeche	Goose Egg	Dinwoody Phosphoria	
PALEOZOIC	PENNSYLVANIAN			Minnelusa Amsden Tyler	Minnelusa	Weber Tensleep Amsden Darwin	Amsden Tyler
				Big Snowy Group Heath Otter Kibbey Madison Group Charles Mission Canyon Lodgepole	Madison Englewood	Madison Mission Canyon Lodgepole	Big Snowy Group Heath Otter Kibbey Madison Group Sun River Charles Mission Canyon Lodgepole
PALEOZOIC	DEVONIAN			Bakken Three Forks Nisku Duperow Souris River Dawson Bay Winnipegosis	Jefferson		Three Forks Nisku Duperow Souris River
		SILURIAN		Interlake	Interlake		
PALEOZOIC	ORDOVICIAN			Stonewall Stony Mountain Red River	Big Horn Winnipeg	Big Horn	Red River
		CAMBRIAN		Winnipeg Deadwood	Deadwood	Gall Gros Flath	Emerson Flathead

Producing Horizon Legend (after Geomap Executive Reference Map, 1983)

**S** = Source Rock



## REGIONAL GEOLOGY

The Fort Berthold Reservation is situated near the deepest part of the Williston Basin (see Fig. FB-2.1 A-A' and associated cross-sections). During the Paleozoic and early part of the Mesozoic, the basin was a stable, cratonic depocenter which received over 15,000' of sediments. Fort Berthold reservation is located within the depocenter, near a major structural feature called the Nesson Anticline, which produces a significant percentage of hydrocarbons within the basin.

Predominantly a carbonate depocenter in the Paleozoic, the basin is also interbedded with clastics and evaporites. The clastic intervals are composed of marine, organic rich shales which are the principal source rocks for the basin. In addition, some of the clastic intervals also include nearshore marine or fluvial sandstone deposits. The carbonate and evaporite units are mainly tidal flat, bioherm/reef or sabhka deposits. Cyclic sedimentation of marine shales, limestones/dolomite, and anhydrites or salt are indicative of the Paleozoic section within the Williston Basin.

Potential reservoir intervals can be formed in the limestone or dolomite via primary or secondary porosity mechanisms. Porosity may be intergranular, vuggy, intercrystalline or fractured or combinations of all types depending on structural position and depositional environment.

### Geologic History - Cambrian and older rocks


Precambrian age supracrustal sedimentary rocks are present in western Montana and extend into Glacier National Park (see Fig. FB-2.1). These rocks are estimated to be from 900 to 1400 million years old. No Precambrian rocks are exposed on the Fort Berthold Reservation.

During Cambrian time, a major seaway existed in western Montana and eastern Idaho (see Figs. FB-2.2 & 2.4). This seaway gradually transgressed from west to east across eastern Montana and the Dakotas. The dominant source of coarse-grained clastics was to the east (from the Sioux Arch) and gradually changed into shales and limestones to the west. Thickness of the Cambrian rocks varies from over 2000 feet in the Montana Disturbed Belt to less than 100 feet along the eastern edge of the Williston Basin. Cambrian sediments buried under the Fort Berthold Reservation are about 300-600 feet thick and composed predominantly of coarse-grained sandstone.


### Geologic History - Ordovician to Triassic

A major depocenter evolved along the eastern edge of the Williston Basin which was a stable, marine shelf area throughout much of the Paleozoic (see Fig. FB-2.3). Ordovician and Silurian rocks were deposited mostly in a shallow tidal flat environment which resulted in alternating cycles of limestone/dolomite, marine shales, and evaporites. By the end of Silurian time, a regional lowstand resulted in a basin-wide unconformity separating Silurian and Devonian rocks. This unconformity influenced the development of vuggy, karsted, carbonate sediments adjacent to this horizon. Present-day thickness of Ordovician and Silurian rocks in the reservation area are 1200 feet and 1000 feet, respectively.


Deposition during Devonian time proceeded much as it had in the Silurian except for the development of highly organic-rich shales within the carbonate intervals. Within the reservation boundaries, Devonian sediments are about 1700 feet thick and include the regional Prairie Salt (500-700'), and the Bakken Shale (70-100'). The Prairie Salt forms a regional seal for the older intervals and has been mobilized/dissolved out of this section near the western edge of the basin (105 degrees longitude). The Bakken Shale is thought to be one of the primary source intervals for Mississippian and younger production.




**Figure FB-2.2.** Generalized time-slice cross-section along A-A'. Line of section along 48 degrees latitude with selected points every 1 degree longitude. Datum is base Ordovician.




**Figure FB-2.4.** Map showing thickness of Cambrian-aged Deadwood and equivalent rocks along with facies information, location of analog fields from Cambrian sediments, location of reservation, and location of regional cross-section A-A' (modified after Peterson, 1987).



**Figure FB-2.1.** Present day structural features of the northern Rocky Mountain region. Includes major fault zones, uplifts, basins, and reservation areas (modified after Peterson, 1987).



**Figure FB-2.3.** Generalized time-slice cross-section A-A'. Triassic through Ordovician. Line of section along 48 degrees latitude with selected points every 1 degree of longitude. Datum is the base of Jurassic, Permian missing (from C.W. O'Melveny, July 1996).



**Figure FB-3.3.** Paleogeographic map of North America during Late Cretaceous time, showing the extent of the Cretaceous seaway (after Rice and Shurr, 1980).

**GEOLOGIC HISTORY (continued)**

**Geologic History - Ordovician to Triassic**

By Mississippian time, the western portion of the Williston Basin was continuously receiving carbonates and evaporites in a shallow, marine shelf environment (see Figure BF-3.1). Most of the producing reservoirs in the basin are from these cyclic marine shales, limestone/dolomite porosity horizons, and evaporitic carbonate sequences. Eventually, the Charles Salt horizon would cover the entire basin and part of eastern and central Montana. By late Mississippian time, deposition of shales and mudstones were mainly confined to the central Williston area and the Big Snowy trough in central Montana. Total thickness of Mississippian rocks within the reservation boundaries is about 2400-2800 feet.

Another major lowstand at the end of the Mississippian time led to widespread erosion and karstification of the underlying carbonate intervals. Pennsylvanian sediments are confined to the center of the Williston basin and central Montana. Pennsylvanian rocks are about 400 feet thick.

Permian deposits are confined to the central Williston basin area and are predominantly sandstone/shale and evaporite sequences. As the Williston basin became filled to base-level, only shallow marine/terrestrial sediments were deposited. This also resulted in numerous unconformities in this horizon. A major unconformity at the end of Permian time has

removed any evidence of these rocks west of longitude 104 degrees. Permian rocks within the reservation are about 500 feet thick. Triassic-aged sediments are also present and of continental origin. Estimated thickness of Triassic rocks across the reservation are about 400-500 feet thick.

**Geologic History - Jurassic to Cretaceous**

A tectonic structural reorganization of the North American continent occurred during Jurassic-Cretaceous time. This resulted in a major change in depocenter position of the Williston basin, shifting from the east to the western side (Figure 4.3). The initial pulses of the Sevier and later Laramide thrusting resulted in dominantly clastic deposition in the Cretaceous Seaway during this time (Figure 4.4).


Thickness of Jurassic rocks across the reservation area are estimated to be about 1200 to 1400 feet thick and are comprised of a complex mixture of nearshore marine, fluvial, and evaporitic deposits. Early Cretaceous-aged continental/fluvial sediments are about 300-400 feet thick. Provenance for these sediments are thought to have been from the southeast in what is present day South Dakota.

The Mowry/Skull Creek Formation is about 400-500 feet thick within the reservation area and was deposited in a transgressive marine sequence which extended from western Montana eastward into the Dakotas; from Texas northward into Canada. Numerous clastic sandstone deposits are present within this sequence and are the result of variations in sea level and clastic influx into the seaway.


During Upper Cretaceous time thrusting and crustal loading from the west had subsided enough to allow the re-establishment of carbonate deposition within the seaway. Extensive chalk deposits of the Greenhorn/Niobrara Formations were deposited as well as thousands of feet of marine carbonate/clastic shale. Upper Cretaceous rocks in the area are more than 2500 feet thick. As the Laramide Orogeny and associated thrusting began to exert influence, nearshore marine and fluvial sandstones began depositing along the shorelines of the seaway.

**Geologic History - Tertiary and Quaternary**

As the orogenic uplifts of the Laramide Orogeny occurred during Late Cretaceous to Tertiary time, older Cretaceous rocks were uplifted and eroded. Only the central portion of the Williston preserved the swamp/peat deposits during the Paleocene and Eocene. Coal deposits of the Fort Union and equivalent rocks are the result. These sediments can be up to 1750 feet thick across the reservation. Alpine glaciers existed in Montana during Quaternary time and extensive glacial lakes and ice sheets covered the reservation area.



**Figure FB-3.2.** Generalized cross-section A-A' - Cretaceous and Older Rocks. Line of cross-section along 48 degrees latitude with selected points every 1 degree of longitude. Datum located at the top of the Cretaceous.



**Figure FB-3.4** Isopach map showing thickness and facies distribution of late Mississippian and Pennsylvanian sediments of the Tyler and Big Snowy Group suite of rocks. Location of Fort Berthold Reservation, any analog fields, and older basement rocks also shown (modified after Peterson, 1987).

## Petroleum Systems

Accumulations of hydrocarbons owe their genesis to several critical factors: generation and migration from source intervals, structural/stratigraphic trapping mechanisms, porous reservoir rocks, and the appropriate timing of formation/generation of these factors. At least four petroleum systems are present within the Williston Basin with numerous underexplored potential hydrocarbon exploration targets. This discussion focuses on the source intervals.

### Source rocks: Generation and Expulsion


At least four source intervals have contributed to the hydrocarbon generation and accumulation patterns within the Williston Basin and all are present in the reservation area.

**Ordovician Winnipeg shale** - A very organic rich shale which exceeds richness values of the Bakken shale in some cases. This interval first entered the oil window in latest Cretaceous/Paleocene time. Peak generation and expulsion occurred between 55-38 mya and some generation continues today. Oils typed to this source are found in the Cedar Creek anticline, eastern Montana, and western North Dakota. However, structures which formed in latest Eocene or after (such as the Nesson Anticline) could not trap the oil migrating from this source. This suggests that much of Winnipeg-sourced oil migrated to the northeastern flank of the Williston Basin where undiscovered oil resource may be present in Ordovician and Silurian strata. This source interval is aerally restricted to the southern and central portions of the basin.


**Bakken Shale** - Known as a world-class source interval, the Bakken has an average of 11.33 wt. % organic carbon. Oil generation was probably initiated about 75 mya with initial expulsion occurring about 70mya (late Cretaceous). Calculations based on pyrolysis data suggest that between 92.3 - 110 billion barrels of oil have been generated from the Bakken. Except for a few fields utilizing the Bakken as the reservoir, significant volumes of Bakken sourced oil have not been discovered to date. Some researchers suggest that most of the expelled Bakken oil is probably lost into the drainage system, where it remains dispersed, at very low saturations (see Figures 2.2 and 2.3 below). Most of the larger structures in the Williston Basin contain mixtures of Lodgepole (Madison) and Bakken oils with the latter at low relative concentrations.

**Lodgepole source interval** - This zone contains predominantly carbonate source horizons with relatively low initial yields; 8 kg HC/t rock. However, large volumes of oil have been discovered typed to this source interval, especially within the Nesson Anticline Petroleum Province. This horizon seems to be geographically restricted to the central and southern portions of the Williston Basin. It appears that migration and trapping efficiencies were much higher in this horizon when compared to the Bakken. This may be due to advantageous timing of structure development relative to expulsion/migration.


**Winnepogosis source interval** - The rich, basal carbonate horizons within this unit (47 kg HC/t rock) are restricted to a starved, Devonian which begins along the northern end of the Nesson anticline and continues north into Canada. This interval charges many of the Waulsortian mounds found in some of the Mississippian-aged sequences.




**Figure FB-4.1** - Location of the Williston Basin with major petroleum provinces and line of section for burial history diagrams indicated. Structural contours are drawn on the base of the Madison Group (after J. Burrus, K. Osadetz, S. Wolf, et al, 1995).



**Figure FB-4.2** - Burial history model with patterns of transformation ratio (TR) at the present and distribution of oil saturations, calculated using a finite model, two-dimensional computer model that simulates oil generation, expulsion, and migration. Generation kinetics determined by experimental data from Williston Basin source rocks. Thermal history model constrained by both present temperature and source rock maturity data. Saturations compared to known patterns of hydrocarbon accumulation within the basin. Saturations between 2-5% represent dispersed oil; saturations above 10% represent oil accumulation or depleted source rocks. Arrows show patterns of active oil migration (after J. Burrus, K. Osadetz, S. Wolf, et al., 1995)



**Figure FB-4.3** - Burial history model with patterns of transformation ratio (TR) at the present and distribution of oil saturations, calculated using a finite model, two-dimensional computer model that simulates oil generation, expulsion, and migration. Generation kinetics determined by experimental data from Williston Basin source rocks. Thermal history model constrained by both present temperature and source rock maturity data. Saturations compared to known patterns of hydrocarbon accumulation within the basin. Saturations between 2-5% represent dispersed oil; saturations above 10% represent oil accumulation or depleted source rocks. Arrows show patterns of active oil migration (after J. Burrus, K. Osadetz, S. Wolf, et al., 1995).



### Play Summary

The diagram and summary charts are coded to the play type number and provide a quick reference to the discovered and undiscovered resource for the reservation area. Also listed are USGS (1996) risk estimates and designations for each of the play types. A qualitative brief review of the summary aspects of each play are also shown.

#### Play Types - Explanation

1. Madison Structure Play
2. Madison Shoreline/Truncation Play
3. Mississippian Lodgepole Wausortian Mound Play
4. Ordovician Red River Play
5. Devonian Nisku/Duperow Play
6. Silurian Interlake Play
7. Pennsylvanian Tyler-Heath Play
8. Pre-Red River Gas Winnipeg/Deadwood Clastics
9. Fractured Bakken Play
10. Shallow Microbial Gas Play

Figure FB-5.1. Schematic diagrams of play types at Blackfeet Reservation


Reservation: Fort Berthold		Total Production ( by province-1996)		Williston Basin		Undiscovered resources and numbers of fields are for Province-wide plays. No attempt has been made to estimate number of undiscovered fields within the Fort Berthold Reservation			
Geologic Province: Central Williston Basin		Oil:		1496 MMBO					
Province Area: Williston Basin (143,000 sq. miles)		Gas:		1735 BCFG					
Reservation Area: 1530 sq. miles (980,000 acres)		NGL:		192 MBNGL					
Play Type	USGS Designation	Description of Play	Oil or Gas	Known Accumulations	Undiscovered Resource (MMBOE)	Play Probability (chance of success)	Drilling depths	Favorable factors	Unfavorable factors
<b>1</b> Madison, structure	3101a	folded structures, primary and secondary porosity in carbonates	Both	878 MMBO 916.5 BCFG 77.9 MMBNGL (numbers include 1, 2, & 3)	<b>Median:</b> 600 MMBO(30 fields @ 20MMBO) <b>Field Size (&gt; 1 MMBOE)</b> 2 MMBO (min), 20 MMBO (median), 5.3 MMBO(mean) <b>No. of undiscovered fields (&gt; 1 MMBOE)</b> 9 (min) 30 (median) 60 (max) 31.9 (mean) numbers include plays 1, 2, & 3	1 high	3,000 - 12,000 ft	1) confirmed play; excellent production within reservation 2) thermally mature source rocks 3) source rocks and reservoir present 4) seismic delineation is useful	1) lack of well control 2) rough topography 3) porosity and facies may be highly variable
<b>2</b> Madison shoreline/truncation play	3101b	Cyclic evaporite/ carbonate sequence, structure/stratigraphic updip pinchout, multiple shoreline cycles	Both	878 MMBO 916.5 BCFG 77.9 MMBNGL (numbers include 1, 2, & 3)	<b>Median</b> of 600 MMBO (30 fields @ 20MMBO) <b>Field Size (&gt;1 MMBOE)</b> 2 MMBO(min) 20 MMBO(median) 5.3 MMBO(mean) <b>No of undiscovered fields (&gt; 1 MMBOE)</b> 9 (min) 30 (median) 60 (max) 31.9 (mean) numbers include plays 1, 2, & 3	1 high	3,000 - 12,000 ft	1) confirmed play; excellent production within reservation 2) thermally mature source rocks 3) source rocks and reservoir present 4) trend extends into reservation 5) mostly shallow drilling depths	1) lack of well control 2) rough topography 3) porosity and facies may be highly variable 4) seismic may not be able to delineate shoreline trends
<b>3</b> Miss. Lodgepole/Wausortian Mound play	3101c	Mound buildups; 'reefs', small but prolific structures; excellent porosity and permeability	Both	878 MMBO 916.5 BCFG 77.9 MMBNGL (numbers include 1, 2, & 3)	<b>Median</b> of 600 MMBO (30 fields @ 20MMBO) <b>Field Size (&gt; 1 MMBOE)</b> 2 MMBO(min) 20 MMBO(median) 5.3 MMBO(mean) <b>No of undiscovered fields (&gt; 1 MMBOE)</b> 9 (min) 30 (median) 60 (max) 31.9 (mean) numbers include plays 1, 2, & 3	1 high	3,000 - 12,000 ft	1) confirmed play; trend probably extends to reservation 2) thermally mature source rocks 3) source rocks and reservoir present 4) seismic may be very useful	1) lack of well control 2) rough topography 3) small areal extent, may be difficult to explore for
<b>4</b> Ordovician Red River Play	3102	Cyclic evaporite/ carbonate sequence, structure/stratigraphic updip pinchouts; multiple shoreline cycles	Both	188.3 MMBO 555.7 BCFG 70.5 MMBNGL	<b>Median</b> of 250 MMBO (25 fields @ 10 MMBO) <b>Field Size (&gt;1 MMBOE)</b> 2 MMBO/10 BCFG(min) 10 MMBO/35 BCFG(median) 2.1 MMBO/11.7 BCFG(mean) <b>No of undiscovered fields (&gt; 1 MMBOE)</b> 5 (min) 25 (median) 50 (max) 26 (mean)	1 high	7,000 - 12,000 ft	1) confirmed play; production within reservation 2) thermally mature source rocks 3) source rocks and reservoir present 4) seismic useful in locating structures	1) lack of well control 2) rough topography 3) possible small exploration targets

Table FB-5.1. Play summary chart

<b>Reservation:</b> Fort Berthold <b>Geologic Province:</b> Central Williston Basin <b>Province Area:</b> Williston Basin (143,000 sq. miles) <b>Reservation Area:</b> 1530 sq. miles (980,000 acres)		<b>Total Production ( by province-1996) Williston Basin</b> <b>Oil:</b> 1496 MMBO <b>Gas:</b> 1735 BCFG <b>NGL:</b> 192 MBNGL			<b>Undiscovered resources and numbers of fields are for Province-wide plays. No attempt has been made to estimate number of undiscovered fields within the Fort Berthold Reservation</b>				
Play Type	USGS Designation	Description of Play	Oil or Gas	Known Accumulations	Undiscovered Resource (MMBOE) Field Size (> 1 MMBOE) min, median, mean	Play Probability (chance of success)	Drilling depths	Favorable factors	Unfavorable factors
<b>5</b> Nisku and Duperow	3103	Cyclic evaporite/carbonate sequences. Structural and stratigraphic pinchouts. Excellent porosity and permeability	Both	160.5 MMBO 159.2 BCFG 12.7 MBNGL	Median 250 MMBO (25 fields @ 10 MMBO) <b>Field Size (&gt; 1MMBOE)</b> 2 MMBO/10 BCFG 10 MMBO/60 BCFG 2.1 MMBO/13.1 BCFG <b>No. of undiscovered fields (&gt; 1 MMBOE)</b> 9 (min) 25 (median) 60 (max) 26.9 (mean)	1 high	8,000 - 12,500 ft	1) confirmed play; production within reservation 2) thermally mature source rocks 3) source rocks and reservoir present 4) seismic delineation is useful	1) lack of well control 2) rough topography
<b>6</b> Silurian Winnipegosis and Interlake	3105	Cyclic evaporite/ carbonate sequence, erosional surfaces. Primary and secondary porosity. Structural and unconformity related trapping mechanisms	Both	55.5 MMBO 180 MMCFG 24.8 MBNGL	Median 225 MMBO (15 fields @ 15 MMBO) <b>Field Size (&gt; 1MMBOE)</b> 3 MMBO/15 MCFG(min) 15 MMBO/90 MMCFG(median) 3.3 MMBO/19.7 MMCF(mean) <b>No. of undiscovered fields (&gt; 1 MMBOE)</b> 5 (min) 15 (median) 25 (max) 15 (mean)	1 mod.-high	8,000 - 12,500 ft	1) confirmed play; production within reservation 2) thermally mature source rocks 3) source rocks and reservoir present 4) seismic useful in prospect delineation	1) lack of well control 2) rough topography
<b>7</b> Post Madison Penn. Tyler/Heath	3106	Fluvial and nearshore sandstones with structural closures. Traps may also occur as discontinuous sandstone lenses.	Both	133.5 MMBO 28.8 BCFG	Median 16 MMBO (8 fields @ 2 MMBO) <b>Field Size (&gt; 1MMBOE)</b> 2 MMBO 10 MMBO 2.1 MMBO <b>No of undiscovered fields (&gt; 1 MMBOE)</b> 4 (min) 8 (median) 15 (max) 8.6 (mean)	1 mod.-high	5,500 - 9,000 ft	1) Thermally mature source rocks 2) source rocks and reservoir present 3) shallow drilling depths	1) No production within reservation 2) rough topography 3) lack of well control 4) depositional area within reservation may be marine instead of shoreline
<b>8</b> Ordovician Pre-Red River Play	3107	Clastic sequences, fluvial and nearshore blanket sandstones. Large, faulted structures	NGL and low BTU gas	no information available	Median 50 BCFG (5 fields @ 10 BCFG) <b>Field Size (&gt; 1MMBOE)</b> 10 BCFG 25 BCFG 13.1 BCFG <b>No. of undiscovered fields (&gt; 1 MMBOE)</b> 1 (min) 5 (median) 20 (max) 7.3 (mean)	1 moderate	10,000 - 16,000 ft	1) confirmed play; production within reservation 2) thermally mature source rocks 3) source rocks and reservoir present 4) seismic useful in locating structures 5) high volume reserves	1) lack of well control 2) rough topography 3) low BTU, contains nitrogen
<b>9</b> Fractured Bakken	3111	Organic rich shale, marine siltstone; fractured; thermally mature oil shale	Both	No information available Oil shows from Sanish sandstones	not estimated 70.3 MMBO/ sq. mile generated hydrocarbons 56.24 MMCFG/ sq. mile generated hydrocarbons Area of play = 8185 sq. miles 7806 sq. miles untested	1 0.2 (20%)	7,500 - 11,100 ft	1) confirmed play; production within reservation 2) thermally mature source rocks 3) source rocks and reservoir present 4) seismic delineation is useful	1) lack of well control 2) rough topography 3) Probable narrow bands of potential fractured reservoir zones
<b>10</b> Niobrara Microbial Gas Play	3113	Niobrara limestone and other shallow reservoirs, self-sourced; porosity decreasing with increasing depth. Large volume accumulations possible	Microbial gas	Only production to date is from Cedar Creek Anticline and Bowdoin dome. These fields are from shallow Eagle Formation sandstones, not Niobrara.	not estimated 180 MMCFG/160 acres (median) 256 MMCFG/160 acres (mean) Area of play = 55,000 sq. miles 29,958 sq. miles untested (mean)	1 0.5 (50%)	500 - 4500 ft	1) large volume play 2) shallow drilling depths 3) accumulations in structural traps, seismic may be useful in delineation of traps	1) lack of well control 2) rough topography 3) reservoir continuity is problematic 4) areal extent may be small

Table BR-6.1. Play type summaries.

 Conventional play type  
 Unconventional/Hypothetical play type



**Blue Buttes Field Parameters**

**Formation:** Mississippian Madison

**Lithology:** Interbedded limestones and dolomites.

**Average depth:** 9200 feet (in reservation area)

**Porosity:** averages 7.7%

**Permeability:** 0.1-8 md, average is 3 md.

**Oil/gas column:** oil 280 feet

**Average net pay:** variable

**Other Formations with shows:** Kibbey sandstone, Kibbey limestone and Charles Formation




Figure FB-7.2. Structure contour map of Blue Buttes Field showing location of cross-section A-A'. Structure on top of Lower Charles Salt.

**PLAY TYPE 1**  
**Folded Structure - Mississippian Carbonate Play**

**General Characteristics** - The Mississippian Madison play is primarily a structural play combined with superimposed facies/porosity changes and pinch-outs. This play is the dominant hydrocarbon producer in the Williston basin. The Madison is subdivided into several producing horizons (see cross-sections below), based on porosity zones. These zones are overlain by evaporite or shale seals. The Charles Salt horizon is a regional evaporite seal which overlies most of the Madison Formation.

Reservoir rocks are generally dolomitized carbonate rocks with either algal, oolitic, crinoidal, or micritic components. Source rocks are thought to be either of Bakken origin or cyclic marine shales within the evaporite-carbonate cycle. Onset of oil generation and migration is modeled to begin in the Late Cretaceous.

**Analog Fields**  
 (\*) denotes fields within Reservation

<b>Antelope*</b> -	39 MMBO	18.9 Mmcf
(includes Bakken, Duperow, and Interlake)		
<b>Blue Buttes</b> -	45 MMBO	28.3 Mmcf
(includes Duperow, Interlake, and Red River)		
<b>Bear Den</b> -	1.4 MMBO	1.5 Mmcf
(Madison, Duperow)		
<b>Croff</b> -	1.7 MMBO	4.0 Mmcf
(Madison, Duperow)		

**Antelope Field Parameters**

**Formation:** Mississippian Madison

**Lithology:** Limestone, brown, dolomitic, fossil fragments, occasional chalky horizons.

**Average depth:** 9100 feet (in reservation area).

**Porosity:** 4.7% gross, intergranular, vuggy

**Permeability:** info. not available

**Oil/gas column:** highly variable

**Average net pay:** variable

**Other shows:** Sanish, Duperow, Interlake.

**Other information:** contains 4.7% H2S





Figure FB-7.3. Antelope field cross-section (after North Dakota Geological Society, 1962).

### Analog Fields

(\*) denotes fields which lie within reservation

**Plaza\*** - 2.6 MMBO, 1.7 Mmcf out of 20 wells, 3-4 MMBO ultimate (Bluell)

**Wabek\*** - 5.1 MMBO, 3.6 Mmcf, out of 18 wells, 6-7 MMBO ultimate (Sherwood)




**Figure FB-8.1.** Sherwood shoreline trend and position of major oil fields ( after Sperr et al, 1993).

### PLAY TYPE 2 Mississippian Shoreline Play

**General characteristics** - This play is an extension of the northeast shelf play which produces from Sherwood and Bluell porosity cycles. In an eastward direction the Mississippian interval subcrops the following formations: Midale, Nesson, Bluell, Sherwood, Mohall, Glenburn, Landa, Wayne, and Lodgepole. Reservoirs are dolomitized carbonates of either algal, oolitic, or bioherm banks along the shoreline trend. The updip seal can either be an evaporite or a shale. Source rocks are likely contained within the Bakken or other marine shales within the evaporite sequence.

#### Plaza Field Parameters


<b>Formation:</b>	Mississippian Mission Canyon, Bluell subinterval
<b>Lithology:</b>	Light brown-brown, peloidal, oolitic pisolitic, intraclastic and composite wackestone-grainstone
<b>Average depth:</b>	7400-7500 feet
<b>Porosity:</b>	intergranular, vugular, intraparticle; 6-16%
<b>Permeability:</b>	no information
<b>Oil/Gas column:</b>	at least 120 feet, no oil/water contact known
<b>Average net pay:</b>	6 feet




**Figure FB-8.2.** Structure map of the Sherwood subinterval - Plaza and Wabek fields (after Sperr et al, 1993).

#### Wabek Field Parameters


<b>Formation:</b>	Mississippian Mission Canyon Sherwood subinterval
<b>Lithology:</b>	Light brown-brown, peloidal, oolitic, pisolitic intraclastic and composite wackestone-grainstone
<b>Average depth:</b>	7300-7500 feet
<b>Porosity:</b>	intergranular, vugular, intraparticle 6-26%, ave.=10%
<b>Permeability:</b>	no information
<b>Oil/Gas column:</b>	at least 100 feet
<b>Average net pay:</b>	26 feet




**Figure FB-8.3.** Wabek Field cross-section showing position of productive interval. Datum is top of Sherwood horizon (after Sperr et al, 1993).




**Figure FB-8.4.** Plaza field cross-section showing position of productive interval. Datum is top of Sherwood horizon (after Sperr et al, 1993).



**WILLISTON BASIN**  
Lodgepole Buildups



**Figure FB-9.1.** Diagrammatic cross-section of Waulsortian Mounds within the Williston Basin, shows facies distribution and general location within the basin (after Burke and Lasemi, 1995).




**Figure FB-9.2.** Generalized isopach map (c.i.=200') of the Lodgepole Formation, Williston Basin in relation to the Fort Berthold Reservation. LB=Little Belt Mountains, B=Bridger Range, BS=Big Snowy Mountains, D=Dickinson Lodgepole Field, S=Saskatchewan (modified from Burke and Lasemi, 1995).

**PLAY TYPE 3**  
**Mississippian Lodgepole**  
**Waulsortian Mounds**


**General Characteristics** - No production has been established within the reservation, however, there is a productive trend in neighboring Stark County. Similar mounds have been found in outcrop in the big Snowy Mountains, Montana.

Waulsortian facies within the Lodgepole formation are lens-like buildups of massive limestone with abundant crinoid and bryozoan fragments. Potential reservoir intervals are boundstones whose framework constituents consist of crinoids, bryozoans, and lesser amounts of mollusks and corals. Inter and intra-particle porosity is the result of leaching and alteration of these particles.




**Figure FB-9.3.** Isopach map of lower Lodgepole at Dickinson Field (after Burke and Lasemi, 1995).

Dickinson Field Lodgepole Parameters	
<b>Formation:</b>	Mississippian Lodgepole
<b>Lithology:</b>	primarily fossiliferous grainstones with minor amounts of dolomite boundstones, packstones
<b>Average depth:</b>	9800 feet
<b>Porosity:</b>	9.4-10% mound core up to 15% in mound flanks
<b>Permeability:</b>	variable, up to 460md
<b>Oil/Gas column:</b>	no information
<b>Average net pay:</b>	at least 50 feet
<b>Other shows:</b>	no information



**Figure FB-9.4.** Generalized Lodgepole section depicting Waulsortian Mound Buildup (after Burke and Lasemi, 1995).



**Figure FB-10.1.** Map showing thickness of Ordovician Red River Formation within the Williston basin and surrounding area, location of analog fields and reservation, and location of regional cross-section A-A' (modified after Peterson, 1987).

**PLAY TYPE 4**  
**Ordovician Red River Play**

**General Characteristics:** This is the second most productive formation in the Williston basin. Reservoirs are dolomite intervals and dolomitic limestones formed from bioclastic mounds and tidal flat deposits. Cyclic deposits of carbonate, evaporite, and organic rich shale provide reservoir, source, and seal. Major accumulations are found on structural noses such as Nesson and Cedar Creek Anticlines. Smaller fields are found in fold structures draped over basement fault blocks, or small carbonate mounds.


The source intervals are thermally mature to overmature at the basin center, and become somewhat immature along the basin flanks. Winnipeg shale and marine shales in the Red River Formation are thought to be the primary source of the reservoir oil. Hydrocarbon generation and migration is estimated to have begun in late Paleozoic time.

**Blue Buttes Field Parameters**


<b>Formation:</b>	Ordovician Red River
<b>Lithology:</b>	black to dark gray dolomite, limestone very fine grained to crystalline occasionally sucrosic texture
<b>Average depth:</b>	-11,300 MSL
<b>Porosity:</b>	9.8%
<b>Permeability:</b>	1.0 md
<b>Oil/Gas column:</b>	unknown
<b>Average net pay thickness:</b>	23 feet
<b>Other shows:</b>	Kibbey Sandstone, Kibbey Limestone Charles Formation
<b>Other information:</b>	Initial IP 564 BOPD, API 58 2928 Mcfgpd-discovery well

**Antelope Field Parameters**


<b>Formation:</b>	Ordovician Red River
<b>Lithology:</b>	black to dark gray limestone/dolomite very fine grained to crystalline Occasionally sucrosic texture
<b>Average depth:</b>	13,480-13,490 feet
<b>Porosity:</b>	12% log density porosity
<b>Permeability:</b>	not known
<b>Oil/Gas column:</b>	no information
<b>Average net pay thickness:</b>	10 feet
<b>Other shows:</b>	Minnelusa and Charles Formations
<b>Cumulative production: (1995)</b>	94 MBO, 1.15 Mmcf API 56.2, IP 113 BC, 1452 Mcfgpd



**Figure FB-10.3.** Red River Structure Map - Blue Buttes Field. Shows trend of Anticline development and production.



**Figure FB-10.2.** Structure contour map of the Red River Fm., Antelope Field. Contours show the general trend of anticline/fold development.



**PLAY TYPE 5**  
**Devonian Nisku-Duperow Play**


**General Characteristics-** This play consists of a carbonate evaporite sequence interbedded with cyclic marine shales. Reservoir rocks are typically dolomite or dolomitized limestone. Source rock for the oil is thought to be from the Bakken interval which is mature-overmature in the central portion of the basin and immature on the flanks. Oil migration and generation are estimated to have begun in early to late Cretaceous time.

Traps are gentle folds and closures related to carbonate bank deposition on paleohighs or shelf areas. These paleostructures are present on regional structural trends such as the Nesson Anticline and Antelope Anticline.

**Analog Fields**  
(\* ) denotes fields which lie within Reservation

<b>Antelope*</b>	39 MMBO, 18.9 Mmcf (includes Bakken, Duperow, and Interlake)
<b>Blue Buttes</b>	45 MMBO, 28.3 Mmcf (includes Duperow, Interlake, and Red River)
<b>Bear Den</b>	1.4 MMBO, 1.5 Mmcf (includes Madison, Duperow)
<b>Croff</b>	1.7 MMBO, 4.0 Mmcf (includes Madison, Duperow)

**Figure FB-11.1.** Map showing thickness of Devonian rocks, limit of Prairie salt, limit of Bakken source rock, location of analog fields and reservation, and location of regional cross-section A-A' (modified after Peterson, 1987).




**Figure FB-11.2.** Bear Den - Devonian Field. Shows position of dolomitic intervals relative to the interbedded evaporite seals. Productive interval indicated in black.

**Bear Den Field Parameters**


<b>Formation:</b>	Devonian Duperow
<b>Lithology:</b>	microcrystalline dolomite with fair microsucrosic porosity
<b>Average depth:</b>	11,300 feet
<b>Porosity:</b>	variable, microsucrosic
<b>Permeability:</b>	not known
<b>Oil/Gas column:</b>	variable
<b>Average net pay thickness:</b>	13 feet
<b>Other info:</b>	no H <sub>2</sub> S

**Antelope Field Parameters**

<b>Formation:</b>	Devonian Duperow
<b>Lithology:</b>	dolomite, brown, finely crystalline, granular to vugular limestone intervals, fossiliferous
<b>Average depth:</b>	10,750 feet
<b>Porosity:</b>	variable, granular, vuggy
<b>Permeability:</b>	not known
<b>Oil/Gas column:</b>	variable
<b>Average net pay thickness:</b>	variable
<b>Other shows:</b>	Madison, Interlake, Sanish
<b>Other information:</b>	No H <sub>2</sub> S



**Figure FB-11.3.** Structure map of Antelope Field. Shows general anticlinal fold trend to the southeast. Inset shows position of Bakken relative to Duperow Formation.



**PLAY TYPE 6  
Pre-Prairie (Winneposis/Interlake Play)**

**General Characteristics** - Regional carbonate units of lower Devonian and Silurian age are overlain by the Prairie Evaporite which acts as a seal rock. Typical reservoirs in the Winneposis are reefs or dolomitized carbonate mounds. Unconformity traps are thought to exist in the Silurian Interlake Formation which can result in dolomitized reefs, minor karsting, and dissolution porosity in tidal deposits.

The Ordovician Red River shales are thought to be the source rocks for this play and are thermally mature within the basin center. Typical traps consist of gentle folds with flexure faulting associated with the regional structure. Stratigraphic traps (either pinch-outs or porosity variations) may exist as well.


**Antelope Field Parameters**

<b>Formation:</b>	Silurian Interlake
<b>Lithology:</b>	dolomite, cream to dark brown possible algal forms, microcrystalline and vugular in part
<b>Average depth:</b>	-9600 feet MSL
<b>Porosity:</b>	variable, granular, vuggy, 7.5%
<b>Permeability:</b>	1.3md
<b>Oil/Gas column:</b>	variable
<b>Average net pay thickness:</b>	variable
<b>Other shows:</b>	Madison, Duperow, Sanish


**Blue Buttes Field Parameters**

<b>Formation:</b>	Silurian Interlake
<b>Lithology:</b>	Dolomite
<b>Average depth:</b>	12,300 feet (-9967 MSL)
<b>Porosity:</b>	12%
<b>Permeability:</b>	not known
<b>Oil/Gas column:</b>	not known
<b>Average net pay thickness:</b>	30 feet


**Figure FB-12.1.** Map showing thickness of Silurian Interlake Formation, facies type, location of analog field and reservation, and location of regional cross-section A-A' (modified after Peterson, 1987).




**Figure FB-12.2.** Example of wireline log through Silurian interval in Blue Buttes Field.




**Figure FB-12.3.** Structure contour map of Interlake interval, Blue Buttes Field. Shows anticlinal nose development with production located somewhat off structure. This indicates a strong stratigraphic component which assists trapping mechanism.




**Figure FB-12.4.** Example of Antelope Field wireline log in the Silurian interval.



**Figure FB-12.5.** Silurian structure map, Antelope Field. Shows anticlinal fold trend to the southeast with production strongly coincident with structure.



**Figure FB-13.1.** Thickness of Upper Mississippian - Lower Pennsylvanian Big Snowy Group interval (Tyler-Heath), location of Fort Berthold Reservation, Dickinson Field (analog), and location of regional cross-section A-A' (modified after Peterson, 1987).




**Figure FB-13.2.** Structure map of Dickinson Field, top of Heath. Complex structural configuration reflects the depositional patterns associated with fluvial, deltaic and nearshore marine environments (after Williston Basin Field Summaries, 1984).

**PLAY TYPE 7**  
**Post Madison Clastics (Tyler-Heath)**


**General Characteristics** - Regional deposition of fluvial, deltaic, and nearshore marine sandstones and carbonates provides the potential reservoirs for this play type. Dark gray to black, organic rich, marine shales of the Tyler are considered to be the main source rock which charge these reservoirs. The shales are thermally mature in the center of the basin and immature along the flanks. Onset of oil generation and migration is thought to have occurred in late Cretaceous to early Tertiary time.

Lateral discontinuity of potential reservoirs in the well-sorted fluvial and nearshore marine sandstones is the norm. In general, areal extent of reservoirs is limited with possible internal porosity and permeability barriers. Overall porosities may be quite good (10-16%). Tyler sandstones are roughly time equivalent to the Morrow sandstones of the mid-continent.



**Figure FB-13.3.** Well log example from Dickinson Field. Upper Mississippian - Lower Pennsylvanian

Dickinson Field Parameters	
<b>Formation:</b>	Pennsylvanian Tyler Mississippian Heath
<b>Lithology:</b>	Interbedded sandstones and shales
<b>Average Depth:</b>	7800 feet
<b>Porosity:</b>	12%
<b>Permeability:</b>	194 md
<b>Oil/Gas Column:</b>	not known
<b>Average net pay:</b>	variable
<b>Other Shows:</b>	shows in deeper Mississippian intervals




**PLAY TYPE 8  
Pre-Red River Gas Play**

**General Characteristics** - Production has been established from Ordovician (Winnipeg) and Cambrian (Deadwood) sandstones. These units are located within the thermally mature or overmature hydrocarbon window of the Williston basin. Both gas and condensate are produced.

Reservoir intervals contain a 'clean' quartz sandstone, silica cement, and enhanced fracture porosity. Source rock is considered to be a marine shale either within the Deadwood or the Winnipeg sandstone. Hydrocarbon generation is thought to have occurred in late Cretaceous to early Tertiary time. Traps are generally asymmetric folds associated with major structural fault zones or hinge lines.


Locations of the fields used as analogs for this play type are noted on the regional facies map. Fort Berthold reservation is bracketed by these fields and in an optimum facies position for possible plays of this type to occur within the boundary of the reservation.

**Figure FB-14.1.** Thickness of Deadwood and equivalent rocks, location of analog fields, location of reservation, and location of regional cross-section A-A' (modified after Peterson, 1987).




**Antelope Field Parameters**

<b>Formation:</b>	Ordovician Winnipeg and Cambrian Deadwood
<b>Lithology:</b>	very fine to fine grained, occasionally medium grained quartz sandstone, occasionally carbonaceous and pyritic
<b>Average Depth:</b>	13,900 feet
<b>Porosity:</b>	12-18% depending upon interval
<b>Permeability:</b>	no information
<b>Oil/Gas column:</b>	no information
<b>Average net pay:</b>	40-50 feet
<b>Other shows:</b>	no information
<b>Other information:</b>	1-32 Brenna-Lacy (1992) completed in Winnipeg-Deadwood. IPF 8BCPD, 5924 MCFGPD. SI for gas. Cumulative production - (1995) 3037 BO, 5.4 MMCF.




**Figure FB-14.3.** Example of Winnipeg-Deadwood formation log signature from Taylor field.



**Figure FB-14.4.** Taylor Field, Winnipeg Structure. Production strongly correlated to major fault with associated anticlinal nose development to the northwest (from Williston Basin Summaries, 1994).


**Taylor Field Parameters**

<b>Formation:</b>	Ordovician Winnipeg and Cambrian Deadwood
<b>Lithology:</b>	Interbedded shales and sandstones Sandstone consists of very fine grained quartz (based on Richardson Field core, Gulf Oil Leviathan 1-21-B)
<b>Average depth:</b>	11,760-11,780 feet
<b>Porosity:</b>	variable, 12-14% density log porosity
<b>Permeability:</b>	no information
<b>Oil/gas column:</b>	no information
<b>Average net pay:</b>	no information
<b>Other shows:</b>	no information
<b>Other information:</b>	Discovery well for Taylor Field, 120 BCPD, 4.54 MMCFPD, 57.9 API. Cumulative production (1995) 128,730 BO, 5.3 MMCF.



**Figure FB-15.1.** Areas of "high" and "low" electrical resistivity in Bakken shales, with subsurface isotherm contours (degrees) and interpreted area of source-rock maturity (after Messiner, 1984).

Fort Berthold reservation is ideally situated for mature Bakken production. The Bakken source interval is thought to have generated over 1 billion barrels of oil but production/migration from the interval is problematic. Mechanisms for emplacement outside the Bakken interval are described below in the west/east cross-section. Production within the Bakken must be concentrated in intervals where fractures (original or induced) can remain open to fluid flow.



**Figure FB-15.2.** Williston Basin with structure contours on the base of Mississippian strata and limit of Bakken Formation (after Webster, 1987).


**PLAY TYPE 9**  
**Bakken Fairway/Sanish Sand Play**

**General Characteristics** - The fractured Bakken Formation can be subdivided into three distinct rock types. The upper and lower zones are black shale with a high organic matter content. The middle zone is a relatively lean organic shale/siltstone. U.S.G.S. analyses of the Bakken indicates that 11.5-12.1 weight percent of the shale is organic carbon. Evidence suggests that the Bakken has generated hundreds of billions of barrels of oil.


The Bakken Fm, where it exists, is thermally mature (see map). It forms a continuously sourced, self-sealed reservoir. Production is

controlled by fractures; matrix porosity and permeability are low. Different fairways are assumed to exist. The areas with the highest potential have elevated thermal maturity, proximity to subcrop, close fracture spacing and proximity to basin flexure hinge lines. Vitrinite reflectance should be greater than 0.9-1.02.

The United States Geological Survey considers Antelope field a special category of Bakken fairway production. The Sanish sand is locally developed, brown, dolomitic, friable, and a slightly argillaceous sandstone with about 6-7% porosity.




**Figure FB-15.3.** Example of log signature from Antelope Field showing Bakken shale interval with sand/silt development




**Figure FB-15.4.** Structure map of the Sanish Pool, Antelope field (from Williston Basin Field Summaries, 1984).

**Antelope Field Parameters**


<b>Formation:</b>	Bakken shale/Sanish sandstone interval
<b>Lithology:</b>	sandstone, dolomitic, brown, friable, slightly argillaceous
<b>Average depth:</b>	10,525 feet
<b>Porosity:</b>	7.4 average
<b>Permeability:</b>	low, changes across structure with the sand/silt content
<b>Oil/Gas column:</b>	no information
<b>Average net pay:</b>	variable
<b>Other formations with shows:</b>	Mission Canyon, Devonian and Winnipegosis
<b>Other information:</b>	Discovery well was Woodward Star-Tribal, sw se 21 T152N R 94 W; 550 BOPD (1953)




**Figure FB-15.5.** Schematic east-west section across the Williston Basin showing source-rock maturity, fluid over-pressure, fracture, migration and hydrocarbon accumulation patterns in the Bakken formation and adjacent units (after Messiner, 1984).



**Figure FB-16.1.** Paleogeographic map of North America during Late Cretaceous time, showing the extent of the Cretaceous seaway (after Rice and Shurr, 1980).



**Figure FB-16.2.** Map showing depth of burial of the Niobrara Formation. Across reservation area porosity could be <35% (after Rice and Shurr, 1980).




**Figure FB-16.3.** General correlation chart of Cretaceous rocks (after Rice and Shurr, 1980).


**PLAY TYPE 10  
Niobrara Microbial Gas Play  
(Low - High Potential)**

**General Characteristics** - Upper Cretaceous Niobrara is a chalk and calcareous shale that covers most of the western interior from Kansas and eastern Colorado into the Dakotas. It is assumed that a Niobrara gas play similar to the eastern Denver Basin (Beecher Island Field, Goodland Field) exists in the southern Williston basin.


Niobrara production in the Denver Basin is considered a self-sourced, continuous extent gas field. Estimated thickness of the Niobrara would be greater than 100 feet, and depth of burial is less than 1000 feet. Area of subcrop or outcrop might affect gas generation. Areal extent of production might be as small as 25 square miles.



**Figure FB-16.5.** Structure map on top of the Niobrara Formation, northwestern Kansas showing a Niobrara gas field (in red). Contour interval is 100 feet. Hypothetical or unconventional play for Fort Berthold reservation (after Lockridge and Sholle, 1978).



**Figure FB-16.4.** Regional distribution of diagenetic and petrophysical facies of the Niobrara. Area within 3000 feet or less of burial should contain chalks with porosity greater than 35%. Areas between 3000 and 4000 feet of burial should average 30-35% porosity (after Rice and Shurr, 1980).



**Figure FB-16.6.** Type logs for Niobrara producing well, Beecher Island area, Kansas Nebraska No. 1-32 Whombie, sec. 32, T2S, R43W (after Lockridge and Sholle, 1978).

## General References

### Fort Berthold Reservation

Anderson, Robert C., 1995, The Oil and Gas Opportunity on Indian Lands-Exploration Policies and Procedures, Bureau of Indian Affairs, Division of Energy and Mineral Resources, General Publication G-95-3, 158 p.

Beeman, William R., et al., 1996, Digital Map Data, Text and Graphical Images in Support of the 1995 Assessment of United States Oil and Gas Resources, United States Geological Survey, Digital Data Series DDS-35, CD ROM.

Charpentier, Ronald R., et al., 1996, Tubular Data, Text, and Graphical Images in Support of the 1995 National Assessment of United States Oil and Gas Resources, United States Geological Survey, Digital Data Series DDS-36, CD ROM.

Gautier, Donald L., et al., 1996, 1995 National Assessment of United States Oil and Gas Resources - Results, Methodology, and Supporting Data, United States Geological Survey Digital Data Series DDS-30 Release 2.

\_\_\_\_\_, et al., 1995, 1995 National Assessment of United States Oil and Gas Resources, Overview of the 1995 National Assessment of Potential Additions to Technically Recoverable Resources of Oil and Gas - Onshore and State Waters of the United States, United States Geological Survey Circular 1118, 20 p.

Mallory, William Wyman, et al., 1972, Geologic Atlas of the Rocky Mountain Region, Rocky Mountain Association of Geologists ,331 p.

Peterson, James A. and MacCary, Lawrence M., 1987, "Regional Stratigraphy and General Petroleum Geology of the U.S. Portion of the Williston Basin and Adjacent Areas", Williston Basin: Anatomy of a Cratonic Oil Province, Rocky Mountain Association of Geologists, pp. 9-43.

Rice, Dudley D. and Shurr, George W., July 1980, "Shallow, Low-Permeability Reservoirs of the Northern Great Plains - Assessment of their Natural Gas Resources", American Association of Petroleum Geologists Bulletin, Volume 64/7, pp. 969-987.

Willette, Donna C., et al., 1996, "Oil and Gas Atlas on Indian Lands", Indian Resources Building Partnerships, Sixth Annual Energy and Minerals Conference, Bureau of Indian Affairs, Division of Energy and Mineral Resources, 10 p.

### Fort Berthold Reservation-Fields and Articles

Anderson, Robert C., 1995, "Fort Berthold Reservation-The Three Affiliated Tribes", The Oil and Gas Opportunity on Indian Lands-Exploration Policies and Procedures, Bureau of Indian Affairs, Division of Energy and Mineral Resources, General Publication G-95-3, pp. 29-42.

Burke, Randolph B. and Zakaria Lasemi, 1995, "A Preliminary Comparison of Waulsortian Mound Facies in the Williston and Illinois Basins", Seventh International Williston Basin Symposium, Montana Geological Society, Billings, Montana, pp. 115-128

Editors, North Dakota Geological Society, 1962, "Antelope Field-Sanish", Oil and Gas Fields of North Dakota-A Symposium , North Dakota Geological Society, Bismark, North Dakota, pp. 21-22.

\_\_\_\_\_, 1962, "Antelope Field-Sanish", Oil and Gas Fields of North Dakota-A Symposium , North Dakota Geological Society, Bismark, North Dakota, pp. 23-24.

\_\_\_\_\_, 1962, "Antelope Field-Duperow", Oil and Gas Fields of North Dakota-A Symposium , North Dakota Geological Society, Bismark, North Dakota, pp. 25-26

\_\_\_\_\_, 1962, "Antelope Field-Interlake", Oil and Gas Fields of North Dakota-A Symposium , North Dakota Geological Society, Bismark, North Dakota, pp. 27-28.

\_\_\_\_\_, 1962, "Antelope Field-Madison", Oil and Gas Fields of North Dakota-A Symposium , North Dakota Geological Society, Bismark, North Dakota, pp. 21-22.

\_\_\_\_\_, 1962, "Bear Den-Madison", Oil and Gas Fields of North Dakota-A Symposium , North Dakota Geological Society, Bismark, North Dakota, pp. 31-32

\_\_\_\_\_, "Bear Den-Devonian", Oil and Gas Fields of North Dakota-A Symposium , North Dakota Geological Society, Bismark, North Dakota, pp. 33-34

\_\_\_\_\_, 1962, "Blue Buttes-Madison", Oil and Gas Fields of North Dakota-A Symposium , North Dakota Geological Society, Bismark, North Dakota, pp. 49-52

\_\_\_\_\_, 1962, "Croff-Madison", Oil and Gas Fields of North Dakota-A Symposium , North Dakota Geological Society, Bismark, North Dakota, pp. 77-78

\_\_\_\_\_, 1962, "Croff-Devonian", Oil and Gas Fields of North Dakota-A Symposium , North Dakota Geological Society, Bismark, North Dakota, pp. 79-80

Editors, Petroleum Information, 1984, Antelope Field-Sanish Pool - McKenzie County, North Dakota", Williston Basin Field Summaries, Petroleum Information.

\_\_\_\_\_, 1984, "Antelope Field-Devonian Pool-McKenzie County, North Dakota", Williston Basin Field Summaries, Petroleum Information.

\_\_\_\_\_, 1984, "Antelope Field-Silurian Pool-McKenzie County, North Dakota", Williston Basin Field Summaries, Petroleum Information.

\_\_\_\_\_, 1984, "Blue Buttes-Silurian Pool-McKenzie County, North Dakota", Williston Basin Field Summaries, Petroleum Information.

\_\_\_\_\_, 1984, "Blue Buttes-Red River Pool-McKenzie County, North Dakota", Williston Basin Field Summaries, Petroleum Information.

\_\_\_\_\_, 1984, "Dickinson Field Heath Pool-Stark County, North Dakota", Williston Basin Field Summaries, Petroleum Information.

\_\_\_\_\_, 1984, "Taylor Field-Winnipeg Pool-Stark County, North Dakota", Williston Basin Field Summaries, Petroleum Information.

Gordon, Ian R., 1995, (Abstract) "Stratigraphy and Structure of an Early Mississippian Waulsortian Bioherm in the Lodgepole Formation, Dickinson Field, North Dakota", Seventh International Williston Basin Symposium, Montana Geological Society, Billings, Montana, pp. 449.

Peterson, James A., 1996, "Williston Basin Province (031)", Tabular Data, Text, and Graphical Images In Support of the 1995 National Assessment of United States Oil and Gas Resources, United States Geological Survey, Digital Data Series DDS-36, CD ROM.

Shurr, George W., et al., 1995, "Tectonic Controls on the Lodgepole Play in Northern Stark County, North Dakota- Perspectives from Surface Studies", Seventh International Williston Basin Symposium, Montana Geological Society, Billings , Montana, pp. 203-208.

Sperr, J.T., et al., 1993, "Wabek and Plaza Fields: Carbonate Shoreline Traps in the Williston Basin of North Dakota", North Dakota Geologic Survey Field Study, No. 1, p. 24.

Webster, Mark and Thomas Van Arsdale; 1995, "Geochemical Microseep Survey of the Plaza and Wabek Productive trends, Mountrail and Ward Counties, North Dakota", Seventh International Williston Basin Symposium, Montana Geological Society, Billings, Montana, pp. 57-65.

### Fort Berthold Reservation - Map References

Executive Reference Map 334, 1985 edition, Extended Area, Northern Rocky Mountains, Geomap Company.

Executive Reference Map 321, 1983 edition, Southern Williston Basin, Geomap Company.

Indian Land Areas, 1992, United States Department of the Interior-Bureau of Indian Affairs.

Clayton, Lee, et al., 1980, Geological Map of North Dakota Survey.

Darton, N.H., et al., 1951, Geologic Map of South Dakota, United States Geological Survey.

Ross, Clyde P., et al., 1958, Geological Map of Montana, Montana Bureau of Mines.

